

Fractional diffusion of cold atoms in optical lattices

Eli Barkai, Physics Department, Bar-Ilan University

Fractional calculus is an old branch of mathematics which deals with fractional order derivatives, e.g., $d^{1/2}/dt^{1/2}$. Recently Davidson's group (Weizmann) has recorded the spatial diffusion of cold atoms in optical lattices, fitting the results to the solution of a fractional diffusion equation

$$\frac{\partial^\beta P(x, t)}{\partial t^\beta} = K_\mu \nabla^\mu P(x, t).$$

Within the semi classical theory of Sisyphus cooling we derive this fractional equation and discuss its meaning and its limitations [1,2]. An asymptotically weak friction force, induced by the laser field, is responsible for the large deviations from normal transport theory (and from Boltzmann-Gibbs equilibrium concepts [3]) at least below a critical value of the depth of the optical lattice.

1. E. Barkai, E. Aghion, and D. Kessler *From the area under the Bessel excursion to anomalous diffusion of cold atoms* **Physical Review X** **4**, 021036 (2014)
2. D. A. Kessler, and E. Barkai *Theory of fractional-Lévy kinetics for cold atoms diffusing in optical lattices* **Phys. Rev. Lett.** **108**, 230602 (2012).
3. A. Dechant, D. A. Kessler and E. Barkai *Deviations from Boltzmann-Gibbs equilibrium in confined optical lattices* arXiv:1412.5402 [cond-mat.stat-mech] (2014).